



# UNION GENERAL ORTHOPEDICS & SPORTS MEDICINE

AFFILIATE OF: UNION GENERAL HEALTH SYSTEM

Jacob B. Stirton, MD

[www.jacobstirtonmd.com](http://www.jacobstirtonmd.com)

706-439-6858

## MULTILIGAMENTOUS KNEE RECONSTRUCTION PHYSICAL THERAPY

### Philosophy

This protocol is to be utilized as a guideline. There will always be individual differences amongst patients regarding progression and tolerance of specific activities. Progression through the protocol will depend on successful accomplishments of set milestones as assessed by the physician and the physical therapist/athletic trainer.

The patient's home exercise program is of utmost importance and should be monitored and emphasized. Initially, patients should be performing their exercises several times a day to regain motion. Due to the importance of regaining early motion, Dr. Stirton's patients are to be seen 3x/week for the first month.

Rehabilitation should create the optimal environment for the natural process of healing to occur. Initially, there should be a strong emphasis on minimizing swelling and pain as well as motion restoration. If a patient's progress is significantly delayed, please contact the physician office to keep them informed.

If you have any questions regarding this protocol, please contact the UGH Orthopedics and Sports Medicine Department at (706) 439-6858.



# UNION GENERAL ORTHOPEDICS & SPORTS MEDICINE

AFFILIATE OF: UNION GENERAL HEALTH SYSTEM

Jacob B. Stirton, MD

[www.jacobstirtonmd.com](http://www.jacobstirtonmd.com)

706-439-6858

## **GENERAL CONSIDERATIONS**

Regardless of the graft choice, patients are to be TTWB with the post-op brace locked in 30 degrees of flexion for 4 weeks for all activities. At 4 weeks, the brace may be locked in full extension and the patient can be WBAT. At 4 weeks the patient will begin PT. They may begin certain WBing exercises IN PT ONLY for patients who have at least 0-90 deg of motion and good quad control.

To minimize joint stress, it is recommended that patients perform closed kinetic chain exercises only in 0-45 degree range with resistance less than full body weight on the injured limb (e.g., leg press, total gym, or bilateral small range squats).

The post-op brace will typically be shortened and unlocked at 8 weeks during the one-month post-operative visit with the surgeon. If there is inclement weather, the patient may need to lock the brace even AFTER the one-month period. An Armour brace will typically be provided at the 3-month visit with the surgeon.

Dr. Stirton's patients will have unrestricted NWBing range of motion UNLESS there was a meniscus repair at the time of surgery in which case ROM is limited from 0-90 degrees for the first 6 weeks.



# UNION GENERAL ORTHOPEDICS & SPORTS MEDICINE

AFFILIATE OF: UNION GENERAL HEALTH SYSTEM

Jacob B. Stirton, MD

[www.jacobstirtonmd.com](http://www.jacobstirtonmd.com)

706-439-6858

## WEEKS 0-4 (Immediate POST-OP PHASE)

### TREATMENT GOALS:

- Minimize swelling/pain.
- **CPM use at home, no PT.**
- ROM: 0-60 deg week 1, 0-90 deg week 2, 0-120 deg week 4.
- Achieve quad activation and improve quad set.

### MODALITIES:

- Cryotherapy and CPM.



# UNION GENERAL ORTHOPEDICS & SPORTS MEDICINE

AFFILIATE OF: UNION GENERAL HEALTH SYSTEM

Jacob B. Stirton, MD

[www.jacobstirtonmd.com](http://www.jacobstirtonmd.com)

706-439-6858

## WEEKS 4-8 (Early Strengthening Phase)

### TREATMENT GOALS:

- Minimize swelling/pain.
- ROM: 0-90 deg week 1, 0-110 deg week 2 (90 if meniscus repaired).
- Gradual progression of WBing with brace locked (unless otherwise noted).
- Achieve quad activation and improve quad set.

Note: It is essential to monitor trunk/core for proper proximal control while doing exercises to avoid substitution.

### PRECAUTIONS:

- No weight bearing ROM exercises yet.
- Gradually increase exercise ROM as dictated by patient response including swelling and soreness response. Articular soreness should be less than 12 hours without medication needed to alleviate symptoms.

### MANUAL THERAPY:

- Patellar mobilizations at every session (patient taught to do at home daily).
- Physiologic stretching flexion/extension.
- Gentle posterior tib-fem mobilizations to start and progress based on patient presentation.

### SUGGESTED EXERCISES:

- Isometrics (quadriceps, gluteals, hamstrings).
- Ankle pumps-> heel raises.
- SLR's (supine with prop under heel as needed, advancing to standing next phase).
- Heel slides (seated or supine); can also do flexion over end of mat table.
- Long sit hamstring stretches (be mindful in patients with hamstring graft).
- AROM knee extension from 90 to 40 degrees.
- Prone hangs working up to 30 minutes/day (3x10 minutes extension stretching daily).
- Prone flexion ROM assisting with opposite LE if needed.
- Prone TKE over ball.
- Prone hip extensions.
- Weight shifting with an active quad set.
- Side lying Abduction series (straight plane ABD, circles, swings) emphasizing neutral spine.
- Trunk stabilization exercises in supine/prone.

Note: High reps, low resistance with focus on proper muscle recruitment.

### MODALITIES:

- NMES quadriceps until no lag sign with SLR, biofeedback with quadriceps exercise, cryotherapy, vasocompression.



# UNION GENERAL ORTHOPEDICS & SPORTS MEDICINE

AFFILIATE OF: UNION GENERAL HEALTH SYSTEM

Jacob B. Stirton, MD

[www.jacobstirtonmd.com](http://www.jacobstirtonmd.com)

706-439-6858

## WEEKS 8-12 (Middle Strengthening Phase)

### TREATMENT GOALS:

- Full flexion and symmetric to opposite side around week 8.
- Normal gait and reciprocal stair management without compensation.
- Fully resolve knee swelling.
- Progression of independent gym/home exercise program.

### PRECAUTIONS:

- To minimize joint stress, it is recommended that patients perform closed kinetic chain exercises only in 0-90 degree range with resistance less than full body weight on the injured limb (e.g., leg press, total gym, or bilateral small range squats).
- No impact activity, pivoting, or twisting.
- No WBing Flexion > 90 deg during exercises.
- Gradually increase exercise ROM and weight as dictated by patient response including swelling and soreness response. Articular soreness should be less than 12 hours without medication needed to alleviate symptoms.

### MANUAL THERAPY:

- Continue with mobilizations as needed to diminish soft tissue and joint restrictions to normal mobility.

### SUGGESTED EXERCISES:

- **Frequency:** Alternate cardiovascular and leg strengthening days with goal of exercising 6 days/week if the patient desires to eventually participate in competitive athletics.
- **Cardiovascular:** Bike, Stair stepper, Elliptical and Retro Treadmill.
- **Strengthening:** Important to focus on quad, hip, and core strengthening.
  - Squats progressing to weighted squats.
  - Progressive step ups/downs (forward, side, back, 4-8" step).
  - ½ Lunges and wall sits in appropriate ROM not to aggravate patellofemoral joint.
  - Single leg balance with opposite leg reaches.
  - One-legged deadlifts.
  - Slide Board.
  - Fast form walking (start in clinic with therapist and progress gradually outside of therapy).
  - Sport cord resisted walking forward/lateral/retro.
  - Swiss ball, planks, and trunk stabilization exercises.
  - 90-40 degree light open chain extension (0-10# ankle weights or TheraBand) in CLINIC ONLY.
  - Static balance exercises using balance cushions, BOSU ball, wobbleboard, half roller, etc.

**MODALITIES:** Cryotherapy, others PRN.



# UNION GENERAL ORTHOPEDICS & SPORTS MEDICINE

AFFILIATE OF: UNION GENERAL HEALTH SYSTEM

Jacob B. Stirton, MD

[www.jacobstirtonmd.com](http://www.jacobstirtonmd.com)

706-439-6858

## **Months 3-4 (Pre-impact training, Progression to Late Strengthening Phase)**

### **SUGGESTED EXERCISES:**

- Continue to advance previous activities as appropriate.
- May start 90-40 degree ROM leg extension machine with high reps, low weight if there is no patellofemoral pain.
- Step to and holds.
- Increase more biofeedback and cueing exercises in regards to knee positioning.
- Advance Closed Chain and Proprioception Exercises.
- Progress with Jump Landing Exercises (give appropriate cues on knee positioning).
- Harvard stepping forward/side to 4-8 inch mat.
- Ladder drills: Starting at 50% speed and gradually progress.
- Alter G or underwater treadmill may be initiated at this time as well.



# UNION GENERAL ORTHOPEDICS & SPORTS MEDICINE

AFFILIATE OF: UNION GENERAL HEALTH SYSTEM

Jacob B. Stirton, MD

[www.jacobstirtonmd.com](http://www.jacobstirtonmd.com)

706-439-6858

## **Months 4-6 (Impact and Late Strengthening Phase)**

### **TREATMENT FOCUS:**

- Maintain Strength and Endurance and Progress to Maximum Strength and Endurance.
- Progress Skill Training.
- Start Jogging and Progress to Running at 5-6 Months.
- Progress Functional/Agility Exercises at 6-7 Months.
- Return to Sport around 9-12 months and scores at least 90% on Functional Testing Compared to Contralateral Side.

### **Running Progression:**

- May begin walk/ slow jog interval progression starting with 1-minute walk and 1-minute jog. Progress increasing jog times by 1 minute and may keep walking time as needed (at least one minute though). When patient can jog 5 minutes straight – progress to a distance-based criteria progression starting at 1 mile. When the patient is running 2 miles without difficulty, then start to add in hills and work on speed.

### **Hop Training:**

- Athletes should be instructed on proper hopping/landing technique. Appropriate cues must be given to the patient to ensure they do not land stiff. They also need cues to avoid LE internal rotation or valgus upon landing.

### **SUGGESTED ADDITIONAL EXERCISES:**

- One-legged hop training.
- More intense two-legged hopping activities.
- Begin  $\frac{3}{4}$  speed sprints if progressed as above on smooth surface.
- Carioca drills (walking->  $\frac{1}{2}$  speed->  $\frac{3}{4}$  speed).
- Figure 8 jogging progression.
- Begin functional sport specific training in controlled environment with trainer or therapist.



# UNION GENERAL ORTHOPEDICS & SPORTS MEDICINE

AFFILIATE OF: UNION GENERAL HEALTH SYSTEM

Jacob B. Stirton, MD

[www.jacobstirtonmd.com](http://www.jacobstirtonmd.com)

706-439-6858

**FUNCTIONAL TESTING:** Please attempt to complete functional testing at 6 months prior to the 6-month doctor's appointment to provide objective data to assist in return to play decision-making. Please make comments on movement quality as well – especially if there are concerns.

- **HOP TESTING:** Include the following hops if possible:
  - Single leg vertical hop.
  - Single leg hops for distance.
  - Single leg 6m timed hop.
  - Single leg triple jump for distance.
  - Single leg triple crossover hops for distance.
- One-minute front step down test at 6-8-inch height. Assess quantity and quality looking for deviations from proper body mechanics.
- Hip/Core strength and control assessment (e.g., planks, one-legged bridge, etc.).
- Sport Specific Testing.

**RETURN TO SPORTS CRITERIA (typically 12-16 months):**

- >90% functional hop testing (limb symmetry index) with good movement quality.
- >90% quadriceps strength.
- >90% One-minute step down test with good movement quality.
- 5/5 hip MMT including hip ER, ABD, and extensors.
- Running/Agility/Sports specific activities are WNL.
- Patient has confidence in leg and does not demonstrate apprehension.