



# UNION GENERAL ORTHOPEDICS & SPORTS MEDICINE

AFFILIATE OF: UNION GENERAL HEALTH SYSTEM

## Post-Op Instructions for Hip Fracture Fixation

Jacob B. Stirton, MD

### DIET

- Begin with clear liquids and light foods (jello, soups, etc).
- Progress to your normal diet if you are not nauseated.

### WOUND CARE

- Maintain your operative dressings.
- It is normal for the hip to bleed and swell following surgery. If blood soaks onto the bandage, do not become alarmed, reinforce with additional dressing.
- Remove surgical dressing on the second day after surgery. Replace dressings with clean 4x4s and tape.
- You can get the incisions wet in the shower (water and soap lightly run over the incisions and pat dry). Simply pad dry afterwards.
- Note that because you cannot place weight on the operative leg that you require the use of a shower chair so that you may safely sit while showering.
- To avoid infection, NO immersion of the operative leg (i.e.: bath or pool) until six weeks after surgery.
- Please do not place any ointments lotions or creams on the incisions.
- Sutures are typically removed between 10-14 days post operatively.

### MEDICATIONS

- A nerve block is typically administered by the anesthesia team unless the patient specifically requests otherwise. This will wear off within 24-36 hours. Patients commonly encounter more pain on the first or second day after surgery when swelling peaks.
- Some patients will require narcotic pain medication for a short period of time – I prescribe oxycodone which can be taken 1-2 tabs every 3 hours as needed for pain.
- To minimize the need for narcotics, I prescribe ibuprofen and Tylenol to be taken three times per day each for the first 2 weeks.
- Common side effects of the pain medication are nausea, drowsiness, and constipation. To decrease the side effects, take the medication with food. We prescribe a stool softener to help prevent constipation but be sure to drink plenty of water as well. If you are having problems with



# UNION GENERAL ORTHOPEDICS & SPORTS MEDICINE

AFFILIATE OF: UNION GENERAL HEALTH SYSTEM

nausea and vomiting, we prescribe an antiemetic, Zofran. If this fails to resolve the issue please contact the office.

- Do not drive a car or operate machinery while taking the narcotic medication.
- Please avoid alcohol use while taking narcotic pain medication.
- For 4 weeks following surgery take one aspirin 325mg tablet daily to lower the risk of developing a blood clot after surgery. Please contact the office should severe calf pain occur or significant swelling of the calf or ankle occur.

## ACTIVITY

- You will either be weight bearing as tolerated or toe touch weight bearing, both with a walker as soon as comfortable immediately after surgery. This depends on the fracture and fixation type.
- **Don't bend your hip past a 90 degree angle.**
- **Don't cross your legs.**
- **Don't twist your hip inwards-keep knees and toes pointed upwards.**
- Do not engage in activities which increase hip pain/swelling (prolonged periods of standing or walking) for the first 7-10 days following surgery.
- Avoid long periods of sitting (without leg elevated) or long distance traveling for 2 weeks.
- NO driving until instructed otherwise by physician.
- May return to sedentary work ONLY 5-7 days after surgery, if pain is tolerable.

## ICE THERAPY

- Icing is very important in the initial post-operative period and should begin immediately after surgery.
- Use ice packs for 20-30 minutes every 2 hours daily for 3 days post-operatively. Avoid frostbite to the skin by not using icepacks for more than 30 minutes at a time.
- You do not need to wake up in the middle of the night to change the icepacks unless you are uncomfortable.

## EXERCISE

- Begin exercises 24 hours after surgery (gait training, stairs, quad sets, straight leg raises, etc.) unless otherwise instructed.
- Complete exercises 3-4 times daily until your first post-operative visit.
- Perform ankle pumps continuously throughout the day to reduce the risk of developing a blood clot in your calf.



# UNION GENERAL ORTHOPEDICS & SPORTS MEDICINE

*AFFILIATE OF: UNION GENERAL HEALTH SYSTEM*

- Formal physical therapy (PT) typically begins the day of surgery. A prescription and protocol will be provided at the time of surgery unless deferred by Dr. Stirton.

## **EMERGENCIES**

\*Contact Dr. Stirton's office at 706-439-6858 if any of the following are present:

- Fever (over 101° F – it is normal to have a low-grade fever (<100°) for the first day or two following surgery) or chills
- Redness around incisions
- Color change in foot or ankle
- Continuous drainage or bleeding from incision (a small amount of drainage is expected)

\*If you have any of the following emergencies that require immediate attention proceed to the nearest emergency room.

- Excessive nausea/vomiting
- Unrelenting pain or calf pain
- Difficulty breathing

## **FOLLOW-UP CARE/QUESTIONS**

- If you do not already have a post-operative appointment scheduled, please contact our scheduler at 706-439-6858 to schedule.
- If you have any further questions please contact Dr. Stirton's office.