



UNION GENERAL ORTHOPEDICS & SPORTS MEDICINE

AFFILIATE OF: UNION GENERAL HEALTH SYSTEM

Jacob B. Stirton, MD

www.jacobstirtonmd.com

706-439-6858

ANKLE/SYNDESMOSIS ORIF PHYSICAL THERAPY

Philosophy

This protocol is to be utilized as a guideline. There will always be individual differences amongst patients regarding progression and tolerance of specific activities. Progression through the protocol will depend on successful accomplishments of set milestones as assessed by the physician and the physical therapist/athletic trainer.

The patient's home exercise program is of utmost importance and should be monitored and emphasized. Initially, patients should be performing their exercises several times a day to regain motion.

Rehabilitation should create the optimal environment for the natural process of healing to occur. Initially, there should be a strong emphasis on minimizing swelling and pain as well as motion restoration. If a patient's progress is significantly delayed, please contact the physician office to keep them informed.

If you have any questions regarding this protocol, please contact the UGH Orthopedics and Sports Medicine Department at (706) 439-6858.

CAUTION: Return to intense activities such as impact loading, jogging, or pivoting and shifting early post-operatively may increase the overall chance of fixation failure. And symptoms of pain, swelling, or instability should be closely monitored and avoided by the patient.



UNION GENERAL ORTHOPEDICS & SPORTS MEDICINE

AFFILIATE OF: UNION GENERAL HEALTH SYSTEM

Jacob B. Stirton, MD

www.jacobstirtonmd.com

706-439-6858

Phase I: Weeks 0- 6

TREATMENT GOALS:

- Control pain and effusion.
- Full ankle ROM.

PRECAUTIONS:

- Non weight bearing for first 6 weeks.
- Splint for first 2 weeks. Then removable boot for next 4 weeks.
- May remove boot to perform NWB exercises and ROM.
- **NWB ROM only** (no limits once out of splint post op week 2)

MANUAL THERAPY:

- Patient **MUST** work on ankle motion at home

SUGGESTED EXERCISES:

- **No loading until 6 wks.**
- Ankle pumps.
- Gastoc/soleus stretch.
- Hamstring/ITB stretch.
- Prone hangs to facilitate extension.
- Heel slides for flexion.
- Quad sets with E-stim.
- SLR in 4 planes, supine/side lying hip circles.
- SAQ, prone knee extensions/TKEs.
- Multi-hip machine in 4 planes.
- Hip flexion-seated.

FOR ADVANCED PATIENT/ATHLETES: UBE, UE weight lifting, core strengthening may be done if it does not load LEs.

MODALITIES:

- Cryotherapy as needed.



UNION GENERAL ORTHOPEDICS & SPORTS MEDICINE

AFFILIATE OF: UNION GENERAL HEALTH SYSTEM

Jacob B. Stirton, MD

www.jacobstirtonmd.com

706-439-6858

Phase II: Weeks 7-12

TREATMENT GOALS:

- Be aware of changes in condition (such as pain and effusion) and modify program as indicated.
- Full ROM.
- Control pain and effusion.
- Full lower extremity strength

PRECAUTIONS:

- Weight bearing as tolerated.
- No Impact.

MANUAL THERAPY:

- Continue stretches as previous.

SUGGESTED EXERCISES:

- Continue previous exercises as indicated.
- Monster walk add variations.
- Heel-toe walking, cone stepping to Dynamic warm-up.
- Leg Press, Total gym, or Reformer.
- Wall squats.
- Lateral step down.
- Stationary bike.
- Mini-squats/squats (0-90).
- Hamstring curl (0-90).
- Leg Press (0-90).
- Lunges-knee not to migrate over toe.
- Begin light circuit training - Stepper, NordicTrack, treadmill, ladder drills.

BALANCE TRAINING:

- Cone walking.
- Mini squat with UE or LE reach (rock around the clock).
- Single leg balance with plyotoss or other challenge.
- Sports cord agility work.
- Wobble board work.

MODALITIES:

- Cryotherapy as needed.



UNION GENERAL ORTHOPEDICS & SPORTS MEDICINE

AFFILIATE OF: UNION GENERAL HEALTH SYSTEM

Jacob B. Stirton, MD

www.jacobstirtonmd.com

706-439-6858

Phase III: Weeks 13+

TREATMENT GOALS:

- Enhance neuromuscular control.
- Perform selected sports specific activity and release per MD to unrestricted sporting activity.
- Achieve maximal strength and endurance.

FUNCTIONAL TRAINING:

- Initiate light plyometric/sports metric type program (as released by MD for impact loading).
- box jumps, level, double-leg, rope jumping, star jumps, hopping.
- Sport specific drills.
- Intensify circuit training - Stepper, elliptical, treadmill, ladder drills, rope jumping, reaction drills.

RUNNING PROGRAM:

- Water walking.
- Swimming (kicking).
- Backward run.
- Return to jogging then running if patient is tolerating plyometrics.

CUTTING PROGRAM:

- Lateral shuffle.
- Carioca, figure 8's.
- LEFTest run.

MODALITIES:

- As needed.

*Advanced weight training and sports specific drills are advised to maintain a higher level of competition.